

Eco 5316 Time Series Econometrics
Lecture 4 Moving Average (MA) processes

MA(q) model

- ▶ let $\{\varepsilon_t\}$ be a white noise time series; process $\{y_t\}$ follows a moving average model of order 1, or MA(1) model, if

$$y_t = c_0 + \varepsilon_t + \theta_1 \varepsilon_{t-1}$$

or, using the lag operator we can write $y_t = c_0 + (1 + \theta_1 L)\varepsilon_t$

- ▶ the bid-ask bounce in stock trading may introduce an MA(1) structure in a return series
- ▶ in general, for $q > 0$ a moving average model of order q , or MA(q) model, is given by

$$y_t = c_0 + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \dots + \theta_q \varepsilon_{t-q}$$

or equivalently using the lag operator $y_t = c_0 + (1 + \theta_1 L + \dots + \theta_q L^q)\varepsilon_t$

Autocorrelation function for MA(q) model

- ▶ if $q = 1$ it can be shown that $\gamma_0 = Var(y_t) = (1 + \theta_1^2)\sigma_a^2$, and also that $\gamma_1 = \theta_1\sigma_a^2$ and $\gamma_l = 0$ for $l \geq 2$; thus

$$\rho_1 = \frac{\theta_1}{1 + \theta_1^2} \quad \rho_l = 0 \text{ for } l > 1 \quad (1)$$

- ▶ theoretical ACF for MA(1) thus cuts off to zero after lag 1
- ▶ if $q = 2$ we have $\gamma_0 = Var(y_t) = (1 + \theta_1^2 + \theta_2^2)\sigma_a^2$, and the theoretical ACF for MA(2) satisfies

$$\rho_1 = \frac{\theta_1 + \theta_1\theta_2}{1 + \theta_1^2 + \theta_2^2} \quad \rho_2 = \frac{\theta_2}{1 + \theta_2^2 + \theta_1^2} \quad \rho_l = 0 \text{ for } l > 2 \quad (2)$$





- ▶ theoretical ACF for MA(1) thus cuts off to zero after lag 2
- ▶ in general, for MA(q) model $\gamma_0 = Var(y_t) = (1 + \theta_1^2 + \dots + \theta_q^2)\sigma_a^2$
- ▶ theoretical ACF for MA(q) model satisfies $\rho_l = 0$ for $l > q$

Autocorrelation function for MA(q) model

- ▶ ACF is thus useful for identifying the order of an MA model in the same way the PACF is useful in identifying the order of an AR model
- ▶ note that MA(q) model is always weakly stationary, unlike AR(p) model which is only weakly stationary if its characteristic roots lie inside the unit circle
- ▶ interactive overview of ACF and PACF for simulated MA(q) models is [here](#)





Autocorrelation function for MA(q) model

MA(1) with $\theta_1 = 0.7$







Autocorrelation function for MA(q) model

MA(1) with $\theta_1 = -0.7$



Autocorrelation function for MA(q) model

MA(2) with $\theta_1 = 0.2$, $\theta_2 = 0.7$



Autocorrelation function for MA(q) model

MA(2) with $\theta_1 = 0.2, \theta_2 = -0.7$

